

Cell free DNA methylation patterns for early detection and management of ovarian and breast cancer

Mukesh Kumar Yadav^{1*}, Santosh Kumar Singh², Manish Singh³,
Shashank Shekhar Mishra⁴, Anurag Singh²,
Jyoti Shankar Tripathi¹ and Yamini Bhusan Tripathi⁵

¹Department of Kayachikitsa, Institute of Medical Sciences, Banaras Hindu University, India

²Centre of Experimental Medicine and Surgery, Institute of Medical Sciences, Banaras Hindu University, India

³Department of Pharmacology, Institute of Medical Sciences, Banaras Hindu University, India

⁴Department of Vikriti Vigyan, Institute of Medical Sciences, Banaras Hindu University, India

⁵Department of Medicinal Chemistry, Institute of Medical Sciences, Banaras Hindu University, India

ABSTRACT

Background: Based on the characteristics of systemic lupus erythematosus-associated pulmonary arterial hypertension (SLE-PAH), Sun et al has put forward a scoring system to distinguish two clinical phenotypes as vasculitic and vasculopathic subtypes[1]. A weighted score ≥ 2 suggested a vasculitic subtype by combining two factors: The time interval between SLE and PAH diagnosis < 2 years and ≥ 2 years were 1 and 0 point; SLE Disease Activity Index (SLEDAI) > 9 , 5-9 and < 5 were 2, 1, 0 point, respectively. While the vasculitic subtype seemed to have poorer prognosis in Sun's research, other study has shown controversial result [2].
Objectives: To find out the prognosis of two distinct clinical phenotypes of SLE-PAH.

Methods: Between 2008 and 2019, a SLE-PAH cohort confirmed by right heart catheterization (RHC) from Guangdong Provincial People's Hospital was included. Other groups of pulmonary hypertension were excluded. Based on the scoring system, patients were divided into vasculitic (weighted score ≥ 2) and vasculopathic subtypes (weighted score < 2). The endpoint was PAH-related mortality. Survival status were confirmed by clinic follow-up data or phone call.

Results: A total of 53 SLE-PAH patients were enrolled. The cases of vasculitic and vasculopathic subtype were 14 and 39, respectively. Ten endpoint events occurred. Eight attributed to PAH and the cause could not be traced in two which were still included in study. The pooled 1-, 3-, 5-year survival rates were 85.7%, 78.6%, 65.5% in vasculitic subtype, and 93.9%, 87.5%, 87.5% in vasculopathic subtype, respectively. Kaplan-Meier analysis showed vasculitic subtype tended to have a poorer prognosis than vasculopathic ($p=0.16$, HR 2.4, 95%CI 0.5-13.8, figure 1).

Bar Chart




Fig.1: Prepare all figures and abstract

CONCLUSION

If authors can accomplish the writing of the 18 paragraphs of text described in this article, they will produce a manuscript that is properly organized, correct in its essentials, and ready for the finishing hand of a seasoned writer and mentor. The secret of getting ahead is getting started. Attributed to Mark Twain (source unknown). Young investigators often finish the data collection and analysis phases of their projects flush with the enthusiasm of finally arriving at an answer, only to find that enthusiasm dwindle as they make their first attempts to write the manuscript. Indeed, the number of abstracts presented at national meetings far exceeds the number of manuscripts that ultimately are published in the medical literature [1]. Such failure to bring good work to publication stems in part from the confusion and perplexity that besets inexperienced writers as they attempt to begin the process of manuscript preparation.

Create a Timetable

It is commonplace that large jobs should be divided into smaller steps with provisional completion dates. Some psychologists recommend conditioned response strategies (defined workplace, timers) to help bring concerted effort to the defined subtask and to keep you from the temptation (and disillusionment) of viewing the project as one monumental and arduous whole. Here is one timetable:

- Session 1: Make notes on the literature, outline template papers
- Session 2: Devise an outline and title for your paper.
- Session 3: Create a rough first draft.
- Sessions 4 and 5: Write revisions one and two.
- Session 6: Write third revision, prepare tables and graphs, then give to coauthors.
- Session 7: Incorporate suggestions from coauthors into the text.
- Session 8: Prepare all figures and abstract.
- Session 9: Proof all changes, check all numbers and units, and review the final product with mentor.

REFERENCES

1. Mentzer JT, DeWitt W, Keebler JS, Min S, Nix NW, Smith CD, et al. DEFINING SUPPLY CHAIN MANAGEMENT. J Bus Logist [Internet]. 2001 Sep 1 [cited 2019 Mar 27];22(2):1–25. Available from: <http://doi.wiley.com/10.1002/j.2158-1592.2001.tb00001.x>
2. Ayers JB. Handbook of supply chain management. St. Lucie/APICS Series on Resource Management, New York.; 2001.
3. Clark M, Barraclough A. MANAGING MEDICINES AND HEALTH PRODUCTS CHAPTER 8 OF HEALTH SYSTEMS IN ACTION [Internet]. Management Sciences for Health; 2010 [cited 2017 Sep 27]. Available from: <http://apps.who.int/medicinedocs/documents/s22123en/s22123en.pdf>
4. Javid Iqbal M, Geer MI, Dar A. Medicines Management in Hospitals: A Supply Chain Perspective A multifaceted Review journal in the field of Pharmacy. Syst Rev Pharm [Internet]. 2018 [cited 2017 Sep 27];8. Available from: http://www.sysrevpharm.org/sites/default/files/10.5530.srp_.2017.1.14.pdf
5. Ministry of Health Nutrition and Indigenous Medicine. Annual Health Bulletin 2016. Medical Statistics Unit Ministry of Health, Nutrition and Indigenous Medicine Sri Lanka; 2018.
6. Avinash Verma Rp. Role of Pharmacist in Supply Chain, Logistics & Storage of Medicine. [Internet]. 2018 [cited 2019 Mar 27]. Available from: <https://www.linkedin.com/pulse/role-pharmacist-supply-chain-logistics-storage-avinash-verma-rph>
7. WHO Regional Office for Africa. Management of Drugs at

Health Centre Level [Internet]. World Health Organization Regional Office for Africa; 2004 [cited 2017 Sep 27]. Available from: <http://apps.who.int/medicinedocs/pdf/s7919e/s7919e.pdf>

8. International Pharmaceutical Federation. Pharmacists in the Supply Chain: The Role of the Medicines Expert in Ensuring Quality and Availability, 2018. International Pharmaceutical Federation (Fip) [Internet]. Essential Medicines and Health Products Information Portal A World Health Organization resource. 2018 [cited 2019 Mar 27]. Available from: <http://apps.who.int/medicinedocs/en/m/abstract/Js23434en/>

9. Pérez-Salazar M del R, Aguilar Lasserre AA, Cedillo-Campos MG, Hernández González JC. The role of knowledge management in supply chain management: A literature review. J Ind Eng Manag [Internet]. 2017 Oct 26 [cited 2019 Jun 28];10(4):711. Available from: <http://www.jiem.org/index.php/jiem/article/view/2144>

*Corresponding author:

Mukesh Kumar Yadav

E mail: mukesh.yadav@bhu.ac.in

Department of Kayachikitsa, Institute of Medical Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh-221005, India.

Tel: 919792353549

Citation: Yadav MK, Singh SK, Singh M, Mishra SS, Singh A, et al. (2018) Neurocognitive Values of *Evolvulus alsinoides* and *Centella asiatica* on Scopolamine Induced Amnesia in Mice. Am J Ethnomed Vol.6 No.1:15.

ANNEX

